

## THE EFFECT OF DIFFERENT SOLVENT POLARITY ON THE PRECIPITATION OF HEAVY ORGANICS FROM A CRUDE OIL DEPOSIT IN THE NIGER DELTA.

**Okoye, I. P, Ofodile, S.E and \*Chukwu, O. C**

*Department of Pure and Industrial Chemistry,  
University of Port Harcourt,  
P.M.B. 5323 Choba,  
Port Harcourt, Nigeria.*

\*Corresponding author. Tel: +2348030480676.  
Email address: [obianujuchukwu2007@yahoo.com](mailto:obianujuchukwu2007@yahoo.com)

Received 11 – 10 – 11

Accepted 27 – 10 – 11

### ABSTRACT

*The precipitation of heavy organics (Saturates, Aromatics, Resins and Asphaltenes) from the Niger delta (Atan crude oil residue) was studied. Two C<sub>4</sub> organic polar solvents (ethyl acetate and butanone) were used to investigate the effect of polarity of the precipitating solvents. The heavy organics precipitated from Atan crude residue were 3.34% and 3.76% for ethyl acetate and butanone respectively. Asphaltenes were precipitated from the heavy organic precipitates using n-C<sub>5</sub> and n-C<sub>6</sub> alkanes. Chromatographic separation and spectroscopic analysis of the deasphalted soluble filtrate were carried out. The results identified the presence of saturates, aromatics and resins in the C<sub>4</sub> heavy organic precipitate.*

**Key words:** Heavy Organics, Precipitates, Saturates, Aromatics, Resins, Asphaltenes

### INTRODUCTION

Crude oil is a naturally occurring flammable liquid consisting of a complex mixture of hydrocarbons and heteroatomic organic compounds of various molecular weights. It is found in geologic formations beneath the earth's surface (Hyne, 2001).

For petroleum fluids, composition and properties vary continuously from the simplest structures to macromolecules. The common problem consists then of identification of the discrete components of the crude, the definition of families or fractions of similar components and understanding the nature and role of interactions between various species. The heavy end is assumed to consist of four major fractions of similar species which are: saturates aromatics, resins and asphaltenes (SARA). At the condition

of the original oil in the reservoir the light and heavy ends are mutually soluble forming one phase so long as certain conditions are maintained in the reservoir (Mansoori *et al.*, 2006) Variations in temperature, pressure or composition may alter this balance and some of the species may segregate and form another liquid or solid phase. Once separated, the complex species may aggregate due to their large size differences, hydrogen bonding or polar group interactions and form larger particles, named flocs (or supramolecules). Flocs could precipitate out of the mixture as solid deposits which in many cases would plug wells, pipelines and processing equipment (Kawanaka *et al.*, 1991).

In SARA, asphaltenes are removed by precipitation with a paraffinic solvent and

deasphalted oil (DAO or maltenes) is separated into saturates, aromatics and resins (SAR) by silica gel chromatography.

Due to the complexity of heavy organics as well as limitations of analytical instruments, the nature of heavy organics (soluble and insoluble) in crude oil, is still not well understood. However, there is some information on the chemical composition of resins and asphaltenes, but little is known about the nature of aggregation process and solvent effects on their solubility, dispersion and their tendency to precipitate from crude oils, which are not well studied and documented. Therefore, butanone and ethyl acetate, two solvents of different polarity were used to investigate the soluble and insoluble heavy organics present in the crude oil. The result of this study would be specifically useful in understanding the behaviour of heavy organics in petroleum fluid.

## **MATERIALS**

### **Experimental**

The atmospheric residue from Atan crude oil used in this study was supplied by the Nigerian National Petroleum Corporation (NNPC), Research and Development Division, Port Harcourt.

## **METHODOLOGY**

### **Precipitation of Heavy Organics Using C<sub>4</sub> Polar Solvents**

Approximately 5g of the residue was added to 100cm<sup>3</sup> of ethyl acetate and butanone respectively. The mixtures were shaken vigorously in a mechanical shaker for 30min and allowed to stand for 48hr before filtration. After filtration, the precipitates contained in the filter paper were oven dried and weighed. The filtrates were concentrated and weighed prior to further precipitation in n-alkanes (n-C<sub>6</sub> and n-C<sub>5</sub>).

### **Dissolution of Precipitates in n-C<sub>6</sub> Alkane**

The solid precipitate was added to 15cm<sup>3</sup> of n-C<sub>6</sub> alkane (non-polar solvents). This was followed by vigorous shaking in a mechanical shaker for 30min, and the solution was allowed to stand for 48hr before filtration.

### **Dissolution of Filtrates in n-C<sub>6</sub> Alkane**

The filtrates were evaporated and solid recovered in each case was added to 30cm<sup>3</sup> of n-C<sub>6</sub> alkane (non-polar solvents). The solution was vigorously shaken for 30min and allowed to stand for 48hr before filtration. After filtration, the precipitates contained in the filter papers were oven dried for 30min and weighed. The filtrates obtained (maltenes) were concentrated for SARA (saturates, aromatics, resins and asphaltenes) analysis and identification using chromatographic and spectroscopic techniques.

### **Re – Precipitation in n-C<sub>5</sub> Alkane**

The solid recovered from n-C<sub>6</sub> filtrate was added to 20cm<sup>3</sup> of n-C<sub>5</sub> alkane (non-polar solvent). The solution was shaken vigorously in a mechanical shaker for 30min and allowed to stand for 48hr before filtration. After filtration, the precipitates contained in the filter paper were oven dried for 30min. The filtrates obtained (deasphalted oil fraction) was finally concentrated and weighed prior to SARA analysis and identification using chromatographic and spectroscopic techniques.

## **RESULTS AND DISCUSSION**

### **Precipitation in C<sub>4</sub> Polar Solvents**

The results of the weight and weight percent of heavy organics precipitated by ethyl acetate and butanone are shown in Table 1. The weight percent obtained (Antan crude) ranged between 3.15% to 3.53% (wt/wt), with mean values of 3.34±0.32% for ethyl acetate, while the percentage precipitate by butanone ranged from 3.58 to 3.93% (wt/wt) with a mean value of 3.76±0.30%. The total solid precipitated by these polar solvents are expected to include saturates, aromatics, resins and asphaltenes

(SARA), in agreement with previous reports (Mansoori, 2007). It is also reported that waxes and heavy aromatics with extensive side chains of n-alkanes are insoluble in polar solvents compared to non-polar n-alkane solvents in which they are soluble (Auflem, 2002).

The difference between the mean values of the heavy organics precipitated by ethyl acetate and that precipitated by butanone is 0.42%, indicating the parity in polarity. Ethyl acetate is slightly less polar than butanone and this difference in polarity is reflected in the comparable total amount of precipitated heavy organics reported in Table 1. As previously stated, waxes and heavy aromatics are known to be sensitive to polarity effect. If this observation was extrapolated to n-alkanes (which are non-polar), the waxes may be soluble even in the lower carbon number n-alkanes. This assertion

may therefore imply that the differences in the solubility of heavy organic precipitates in organic solvents of same number of carbon atoms but different functional groups may be as a result of the presence of different groups of compounds in the heavy organic precipitates. (Shehu *et al.*, 1991). Mansoori, 2007 showed that the kind and amount of depositions of heavy organics from petroleum fluids vary, depending on the hydrocarbons present in the oil and the relative amounts of each family of heavy organics.

The weight percent of heavy organics (soluble filtrate) in ethyl acetate and butanone solvents are shown in Table 2. The results in Table 2 show that 88.55% and 91.42% (wt/wt) of the Antan crude oil residue were soluble in ethyl acetate and butanone respectively.

**Table 1: Comparison of heavy organics precipitated using ethyl acetate and butanone polar solvents**

Test Crude Oil	Ethyl acetate			Butanone		
	Wt. of residue (g)	Wt. of HO (g)	Wt. % of HO	Wt. of residue (g)	Wt. of HO (g)	Wt. % of HO
Antan	5.09	0.17	3.34±0.32	5.06	0.19	3.76±0.30

**Table 2: Heavy organics soluble in ethyl acetate and butanone polar solvents (Filtrate)**

Test Crude Oil	Ethyl acetate			Butanone		
	Wt. of residue (g)	Wt. of soluble (g)	Wt. % of soluble (g)	Wt. of residue (g)	Wt. of soluble (g)	Wt. % of soluble
Antan	5.09	4.65	91.42±1.56	5.06	4.48	88.55±1.65

**Re – Precipitation in n-C<sub>6</sub> Alkane**

The results presented in Table 3 show the content of asphaltenes precipitated using n-C<sub>6</sub> alkane. About 37.23% and 21.09% (wt/wt) of asphaltenes were part of ethyl acetate and butanone heavy organic precipitates respectively, and which were insoluble in n-C<sub>6</sub> alkane. This indicates that the heavy organics precipitated from crude oil residue (using polar solvents) contained n-C<sub>6</sub> asphaltenes. It was also observed that n-C<sub>6</sub> heavy organics (insoluble) was hard and dark coloured. This n-C<sub>6</sub> precipitates were thought to be principally asphaltenes because waxes, aromatics and resins were soluble in n-C<sub>6</sub> alkane except where co-precipitation occurred (Rossamdana, 1996; Koka *et al.*, 1993).

The amount of n-C<sub>6</sub>-asphaltene precipitation from ethylacetate (0.21%) and butanone (0.22%) soluble filtrate are also indicated in Table 3. The results indicated that asphaltene precipitated under this condition are not significant, suggesting that asphaltenes are not soluble in these C<sub>4</sub> organic polar solvents. Previous report by Vazquez and Mansoori (2000) showed that the fraction of heavy organics insoluble in n-pentane is larger than that insoluble in n-heptane and contained wider range of molecules, including the fraction of n-pentane insoluble that was soluble in n-heptane. The n-C<sub>6</sub> soluble filtrates obtained were brown in colour. These soluble filtrates were the maltenes used for uv – spectroscopy and silica gel chromatographic qualitative identification of saturates, aromatics and resins.

**Table 3: n-C<sub>6</sub> Asphaltenes content of heavy organics (HO) precipitates and soluble filtrates obtained in C<sub>4</sub> polar solvents (ethyl acetate and butanone)**

	<b>Precipitate (via Ethyl acetate)</b>	<b>Precipitate (via Butanone)</b>	<b>Filtrate (via Ethyl acetate)</b>	<b>Filtrate (via Butanone)</b>
<b>Total Heavy organics (SARA/g)</b>	0.17	0.19	4.65	4.48
<b>n-C<sub>6</sub> Asphaltene(g)</b>	0.06	0.04	0.01	0.01
<b>n-C<sub>6</sub> Asphaltene (%)</b>	35.29±3.06	21.05±1.87	0.21±0.01	0.22±0.01
<b>A/SAR (asphaltene/saturates, aromatics and resins)</b>	0.353	0.211	0.0022	0.0022

**Table 4: n-C<sub>5</sub> Asphaltene precipitated from Antan Crude by n-C<sub>5</sub> via n-C<sub>6</sub> filtrate**

Test	Ethyl acetate			Butanone		
	Wt. of filtrate solid (g)	Wt. of asphaltene (g)	Wt. % of asphaltene	Wt. of filtrate solid (g)	Wt. of asphaltene (g)	Wt. % of asphaltenes
1	0.09	0.03	33.33	0.13	0.04	30.77
2	0.09	0.03	33.33	0.14	0.05	35.71
3	0.09	0.03	33.33	0.15	0.05	33.33
Mean	0.09	0.03	33.33±0.00	0.14	0.05	33.27.07±4.17

### Re – Precipitation in n-C<sub>5</sub> Alkane

The results of weight percent of n-C<sub>5</sub> asphaltenes precipitated from n-C<sub>6</sub> filtrates are presented in Table 4. It was observed that 33.27 to 33.33% (wt/wt) of the solid recovered from n-C<sub>6</sub> soluble filtrates containing C<sub>4</sub> heavy organic precipitates were insoluble in n-C<sub>5</sub> solvent. Thus, n-C<sub>6</sub> soluble filtrates contained n-C<sub>5</sub> insoluble asphaltenes. It also showed that both ethyl acetate and butanone precipitates include n-C<sub>5</sub> insoluble precipitates (or n-C<sub>5</sub> asphaltenes). In essence, the dissolution and re-precipitation using n-C<sub>5</sub> alkane is to investigate the effectiveness of n-C<sub>6</sub> alkane as a heavy organic precipitant. Asphaltenes were part of the hydrocarbon types as defined by SARA responsible for the differences in the quantity of heavy organics precipitated by organic solvents of different carbon chain lengths and polarity. It was also observed that the n-C<sub>5</sub> insoluble precipitates obtained were hard and dark in

colour and the precipitates were n-C<sub>5</sub> asphaltenes in accordance with similar report by Vazquez and Mansoori, 2000. The mean values of n-C<sub>6</sub> and n-C<sub>5</sub> asphaltenes is 37.23±3.06% (for ethyl acetate); 21.09±1.87% (for butanone) and 33.33±0.00% (for ethyl acetate); 33.27±4.17% (for butanone). The total n-C<sub>6</sub> and n-C<sub>5</sub> asphaltenes obtained from ethyl acetate heavy organic precipitate is 1.93 while 1.78 was obtained from butanone heavy organic precipitate (Table 5). The differences between the total asphaltenes is negligible (0.15) compared to 0.42 in the total heavy organics precipitated by C<sub>4</sub> solvents. This suggests that the other compound types – saturates, resins and aromatics were a large part of the precipitates by C<sub>4</sub> solvents. The absorbances (340 – 550nm) of the maltenes and deasphalted oil, after fractionation showed the presence of saturates, aromatics and resins.

**Table 5: Comparism of SARA content in C<sub>4</sub> HO precipitates**

C <sub>4</sub> HO PPT (%)		n-C <sub>6</sub> PPT From C <sub>4</sub> HO (%)		n-C <sub>6</sub> Asphaltene From C <sub>4</sub> HO (%)		n-C <sub>5</sub> PPT From C <sub>4</sub> HO (%)		n-C <sub>5</sub> Asphaltene From C <sub>4</sub> HO (%)	
Ethyl acetate	Butanone	Ethyl acetate	Butanone	Ethyl acetate	Butanone	Ethyl acetate	Butanone	Ethyl acetate	Butanone
3.34	3.76	37.23	21.09	1.24	0.79	33.33	33.27	0.69	0.99

### CONCLUSION

This investigation showed that butanone being more polar, produced greater quantity of heavy organic precipitates than ethyl acetate because waxes and heavy aromatics with extensive side chains were insoluble in polar solvents. It is noted that n-C<sub>6</sub> alkane failed to precipitate all n-C<sub>5</sub>-asphaltenes. Since asphaltenes were defined as the component of crude oil which are insoluble in n-pentane, the n-C<sub>5</sub> filtrates were regarded as the deasphalted oil because the asphaltenes have been removed. It contains the remaining three fractions; saturates aromatics

and resins, which were separated and identified using silica-gel chromatographic and spectroscopic techniques. Quantitative analysis of the saturates, aromatics, resins and asphaltenes (SARA) can be carried out using high performance liquid chromatography (hplc) and/or gel permeation chromatography (gpc).

### REFERENCES

- Andersen, S.I., Speight, J. G. (2001). Coke Generation and Asphaltene Adsorption. *J.PetSci. Tech*; 19, 1 – 34

- Auflem, I. H; (2002). Influence of Asphaltene Aggregation and Pressure on Crude Oil Emulsion stability, PhD thesis, Department of Chemical Engineering Norwegian University of Science & Technology, Trondheim
- Branco, V. A. M., Mansoori, G. A., De Almeida Xavier, L. C., Park, S. J., Manafi, H. (2000). Asphaltene Flocculation and Collapse from Petroleum Fluids. *J. Pet. Sci. Eng.*, 32, 217 – 230.
- Hu, Y.F., Li, S., Liu, N., Chu, Y. P., Park, S. J., Mansoori, G. A., Guo, T. M. (2004). Measurement and Corresponding States Modelling of Asphaltenes Precipitation. *J. Pet. Sci. Eng.*, 41, (1 – 3), 199 – 212.
- Hunt, J. M. (1996). Petroleum Geochemistry and Geology. Freeman, New York. pp.9
- Hyne, N. J. (2001). Non-technical Guide to Petroleum Geology, Exploration, Drilling and Production. Penwell. pp 4.
- Jewell, D. M., Albaugh, E. W., Davis, B. E. and Ruberto, R. G. (1974). Integration of Chromatographic and Spectroscopic Techniques for the Characterization of Residual Oils. *J. Ind. Eng. Chem, Fundam.*, 13, (3), 278 – 282.
- Kawanaka, S., Leontaritis, K. J., Park, S. J., Mansoori, G. A. (1989). Thermodynamic and Colloidal Models Of Asphaltene Flocculation. Oil Field Chemistry, ACS Symposium Series, (396), Washington, D. C., Chapter 24.
- Kawanaka, S., Park, S. J., and Mansoori, G. A. (1991) Organic Deposition from Reservoir Fluids: A Thermodynamic Predictive Technique. *SPE Reservoir Eng. J.*, 185 – 192.
- Kokal, S.I.; Majmanm, J; Sayegh, S. G. And George A. E. (1993). Measurement and Correlation of Asphaltene Precipitation from Heavy Oils by Gas Injection. *J. Can. Pet. Tech*; 31, 24.
- Leontaritis, K. J. and Mansoori G. A. (1989). Fast Crude Oil Heavy Component Characterization using Combination of ASTM, HPLC, and GPC Methods. *J. Pet. Sci. & Tech.*, 2, 1 – 12.
- Mansoori G. A. (1996). Asphaltene, Resins and Wax Deposition From Petroleum Fluids. *Arabian J. Sci. & Eng.* 21, (48), 707.
- Mansoori, G. A. (1997). Modelling of Asphaltene and other Heavy Organic Deposition. *Pet. Sci. & Eng. J*; 17, 101 – 111.
- Mansoori, G. A. (2007). Diamondoids; Their Role in Petroleum and Natural Gas Production Fouling and the Emerging Fields of Biotechnology and Nanotechnology (BioNano Technology). Department of Bio and Chemical Engineering, University of Illinois at Chicago, United States.
- Mansoori, G. A. (2007). Depositions from Petroleum Fluids. Department of Bio and Chemical Engineering, University of Illinois at Chicago, United States.
- Mansoori, G. A., Vazquez, D. and Shariaty-Niassar, M. (2006). Polydispersity of Heavy Organics in Crude Oil and their Role in Oil Well Fouling. Department of Bio and Chemical Engineering, University of Illinois at Chicago, United States.
- Mohamed, R. S. and Ramos, A. C. S. (1999). Aggregation Behaviour of two Asphaltenic Fractions in Aromatics Solvents. *J. Energy & Fuels*, 13, 323 – 327.
- Okoye, I. P. (2009). Fundamentals of Petroleum and Hydrocarbon Chemistry. Allwell press, Port Harcourt, Nigeria. 1st ed. pp. 68 – 78.
- Pacheco-Sanchez, J. H. and Mansoori, G. A. (1989). Prediction of the Phase Behavior of Asphaltene Miscelle/ Aromatic

- Hydrocarbon Systems. *J. Petrol Sci. & Technology*, 16, (3 – 4), 377 – 394
- Park, S. J., Escobedo, J., Mansoori, G. A. (1993). In: Yen, T. F., Chilingarian, G. V. (eds), *Asphaltenes and Asphalts*, Elsevier, The Netherlands.
- Park, S. J., Mansoori, G. A. (1988a). Organic Deposition from Heavy Petroleum Crudes (a FRACTAL Aggregation Theory Approach). Proceed UNIAR/UNDP 4<sup>th</sup> Intl. Conference on Heavy Crudes and Tar Sands, Edmonton, Alberta, Canada.
- Park, S. J. and Mansoori, G. A. (1988b). Aggregation and Deposition of Heavy Organic in Petroleum Crudes. *Intl. J. Energy Sources*, 10, 109 – 125.
- Rassamdana, H; and Sahimi, M; (1996). Asphaltene Flocculation and Deposition: Onset of Precipitation, *AIChE, J*; 42, 3318.
- Shehu, E. Y., Mullins, O. C. (1995). *Asphaltenes: Fundamentals and Application*. Plenum press, New York. pp. 30 – 33, 73 - 76.
- Shehu, E. Y; Store, D. A; De Tar, M. M. (1991). Asphaltenes in Polar Solvents. *J. Non-crystalline Solids*, 341, 131 – 133.
- Vazquez D. and Mansoori (2000). Analysis of Heavy Organic Deposits. *Soc.Pet. Eng. J.*, 26, (1 – 4), 49 – 56
- Wang, J. (2000). Heavy Organic Deposition within Reservoir Rocks. Ph.D. Thesis, New Mexico Institute Of Mining And Technology, Socoro.
- Yen, T. F. (1974). Structures of Petroleum Asphaltenes and its Significance. *Energy Sources*, 1, (4), 447 -463.