

SYNTHESIS, CHARACTERIZATION AND ANTIBACTERIAL ACTIVITY OF Fe(II) MIXED LIGANDS COMPLEX OF SUCCINIC ACID AND HEXAMETHYLENETETRAMINE

J. I. Mbonu¹ and M. Ariemudwigho

Department of Chemistry, College of Science,
Federal University of Petroleum Resources, Effurun,
Delta State, Nigeria

¹Correspondence author e-mail: idongesitmbonu@yahoo.com

Received: 05-03-15

Accepted: 02-04-15

ABSTRACT

An Fe(II) complex has been synthesized from iron(II) sulphate, succinic acid and hexamethylenetetramine. The nature of bonding and the structural features of the complex have been deduced from elemental analysis, molar conductance, pH measurements, IR, UV-Vis spectral studies. Spectroscopic and analytical studies reveal octahedral geometry for this complex. The electronic spectrum of iron(II) complex in DMSO was recorded, and its salient features are reported. The activity studies data indicate that the complex is more potent antibacterial agent than the ligands and ciprofloxacin drug against the tested pathogenic bacteria strains. The synthesized compound can serve as potential photoactive materials as indicated from their characteristic absorption in the ultraviolet region.

Key Words: Succinic acid, hexamethylenetetramine, antibacterial activity.

INTRODUCTION

Many microorganisms, which cause damage to human health, exhibit drug resistance due to improper and abusive use of antimicrobial agents called antibiotics. Antimicrobial resistance has a significant public health and economic impact. Thus, there is a need for measures to reduce this problem. Emphasis on health care costs has provoked a renewed interest in the design and development of novel and cost effective antimicrobial agents with increased biological activity against these resistant strains [1-8]. Research has shown that design and synthesis of transition metal complexes derived from mixed base ligands might be a possible way to obtain materials endowed with antiviral, antibacterial and antifungal properties [3]. Broad empirical screening of chemical compounds for

antimicrobial activity represents an alternative strategy for the development of new antimicrobials complexes with enhanced microbial activities, selective toxicity and chemical stability [4, 9]. The following bacteria *Staphylococcus aureus*, *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Enterococcus species*, *Pseudomonas species* and *Klebsiella species* are associated with a high incidence of infections that are resistant to treatment with antibiotics [2]. Our interest in synthesis of metal complex with antibacterial activities has led to the report herein the synthesis, spectroscopic studies and biological activities of mixed ligands complex of iron(II) with succinic acid and hexamethylenetetramine which have donor and acceptor property due to electron-withdrawing and releasing substituents (OH

and NH_2 groups). We are also interested to examine the optical properties of this compound.

MATERIALS AND METHODS

All the chemicals and solvents used in this study viz, hexamethylenetetramine (HMTA, $\text{C}_6\text{H}_{12}\text{N}_4$), succinic acid ($\text{C}_4\text{H}_6\text{O}_4$), iron(II) sulphate heptahydrate salts ($\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$), ethanol, methanol, ammonia were all obtained from Sigma Aldrich and were of analytical grade. The solvents were used without further purification. Melting point determination was carried out using a Thomas Hoover capillary melting point apparatus. Infrared spectra of the ligands and complex were recorded using pressed KBr discs in the range ($4000\text{-}450\text{ cm}^{-1}$) on Perkin-Elmer FT-IR spectrometer 1.00 versions (Wattman) and the Electronic spectrum was measured in the range $200\text{-}700\text{ nm}$ at room temperature with Perkin Elmer spectrum Bk UV-Vis spectrophotometer.

Synthesis of the Iron(II) Complex:

To a stirred solution of hexamethylenetetramine (5 mmol, 0.701 g) in 10 ml methanol, 5 ml aqueous solution of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (5 mmol, 1.390 g) was added dropwise with stirring at $20\text{ }^\circ\text{C}$ for 10 minutes. Then a 5 ml methanol solution of succinic acid (5 mmol, 0.591 g) was added dropwise with stirring to the mixture. The resulting mixture was stirred for three hours. The reddishbrown complex formed was filtered, washed with methanol and dried under vacuum. On the fifth day, brown crystals were observed to form from the precipitate.

Biological Activity

Antibacterial Screening: All the species used for this test were derived from stock cultures obtained from Environmental

Science Laboratory, Federal University of Petroleum Resources, Effurun, Delta State, Nigeria. Antibacterial activity was tested against *Staphylococcus aureus*, *Staphylococcus aubeus* and *Escherichia coli* using the paper disc plate method. The antibacterial activities of the ligands, and metal complex were evaluated using disc-diffusion method. Nutrient Agar was employed as bacterial growth medium. Antibacterial agents ($100\text{ }\mu\text{g}$) of the complex were loaded on the discs by soaking in test solution, and the discs were placed on the surface of the sterile nutrient agar medium with aid of the sterile forceps and the plates were incubated at $37\text{ }^\circ\text{C}$ for 24 h. Antibacterial activity was evaluated by measuring the diameter of growth inhibition zone (IZ) in mm around the discs using Zone Reader. Compound was considered active when the IZ was greater than 6 mm. Antibacterial activity studies were performed in triplicate, and the average was taken as the final reading. Hexamethylenetetramine, succinic acid and ciprofloxacin were recorded under similar conditions [3].

RESULTS

The physical properties of the complex and ligands listed in Table 1 shows that the Fe(II) complex was isolated as brown crystals. The iron(II) complex decomposed on melting within a temperature range of $160\text{ - }163\text{ }^\circ\text{C}$. The variation in the color with increasing temperature could be due to changes in the crystal structure. The Fe(II) complex is air stable and insoluble in water and partially soluble in ethanol (Table 2). The complex and its ligands were generally soluble in methanol and ammonia.

Table 1: Physical properties of ligands and Fe (II) complex

Ligand/complex	Colour	Physical state	yield (%)	Melting point (°C)
(CH ₂) ₆ N ₄	colorless	crystalline	-	200
C ₄ H ₆ O ₄	colorless	crystalline		186
FeSO ₄ .7H ₂ O	colorless	crystalline	-	100
Fe(II) complex	brown	crystalline	68.5	160-163

Table 2: Solubility of the ligands and Fe(II) complex in water and some solvents

Compound	Solvents			
	Water	Ethanol	Methanol	Ammonia
(CH ₂) ₆ N ₄	Soluble	Insoluble	Insoluble	Soluble
C ₄ H ₆ O ₄	Insoluble	Insoluble	Insoluble	Soluble
FeSO ₄ .7H ₂ O	Insoluble	Soluble	Soluble	Soluble
Fe(II) complex	Insoluble	Partially soluble	soluble	soluble

The elemental composition was determined from energy dispersive X- ray analysis (EDX) spectrum to be calculated: C 15.42,

H 15.38, N 10.22, O 16.26, Fe 42.72 %; found: C 15.98, H 15.56, N 9.28, O 16.34, Fe 42.84 %.




Figure I: EDX spectrum of Fe(II) complex

The infrared spectrum of the complex (Fig.2) was compared with that of the free ligands and qualitative differences between the spectra used in diagnosing the different vibrational frequencies of the metal and ligands in the complex. The Fe(II)

vibrational bands were observed at 2991, 2053, 1445, 977, 697 cm^{-1} assigned to $\nu_{\text{as}}(\text{CH}_2)$, $\nu_{\text{s}}(\text{CH}_2)$, $\nu(\text{C-N})$, $\delta(\text{CH}_2)$, $\nu(\text{M-O})$ vibrational modes, respectively. Similar vibrational frequencies are recorded in the literature for similar compounds [11, 12].




Figure 2: IR spectrum of Fe (II) complex

The electronic spectral data of the complex recorded in dimethylsulfoxide in the range 200–700 nm using a quartz cuvette of 1 cm path length presented in Table 3 confirms

the formation of the iron(II) complex. The UV–vis spectral data values of the complex have shifts frequency when compared with the free ligands.

Table 3: Band positions (nm) and proposed transitions for Fe(II) complex

Complex	Band position	Assignment
Fe (II) complex	203	${}^4T_{1g}(F)$
	253	${}^4T_{1g}(P)$
	269	${}^4T_{1g}(F)$ ${}^4A_{2g}$

From the antibacterial profile presented in Table 4; Fe (II) complex displays a significant inhibitory effect on the growth of tested bacteria species (Fig.3).

**Figure 3: Bacteria inhibition zone with Fe (II) complex****Table 4: Antibacterial activity of ligands, complex and ciprofloxacin-diameter of zone of inhibition (mm).**

Tested compounds	<i>Staphylococcus aureus</i> (G+)	<i>Staphylococcus aureus</i> (G+)	<i>Escherichia coli</i> (G-)
1	0.0	0.0	0.0
2	0.0	0.0	3.6
3	48.1	48.6	20.2
4	23.7	32.2	14.5

Key: G+ = gram positive bacteria, G- = gram negative bacteria, IZ = inhibition zone, 1 = $(CH_2)_4N_4$, 2 = $C_4H_6O_4$, 3 = Fe(II) complex, 4 = Ciprofloxacin drug.

DISCUSSION

The results obtained from solubility tests showed that the complex is soluble in coordinating solvents. The molar conductance measurement of complex was $2.75 \Omega^{-1} \text{cm}^2 \text{mol}^{-1}$ indicating that it possesses a non electrolytic nature. These observations are in agreement with octahedral Fe(II) complexes reported in the literature [10,11].

The experimentally found values for the elemental composition were very close with the calculated values [1]. These data are consistent with the general formulation as 1:1:1, mixed metal-ligands stoichiometry.

The electronic absorption spectrum of the iron (II) complex reveals three bands at 203, 253 and 269 cm^{-1} assigned to ${}^4\text{T}_{1g}(\text{F})$, ${}^4\text{T}_{1g}(\text{P})$ and ${}^4\text{T}_{1g}(\text{F}) \rightarrow {}^4\text{A}_{2g}$ transitions which arise from an octahedral structure [10, 11].

The hexamethylenetetramine ligand (HMTA), was found to be active against only one of the 3 tested bacterial strains (*S. typhi*); whereas the succinic acid has relatively low antibacterial activity [8].

The mixed ligands complex of Fe(II) with succinic acid and hexamethylenetetramine was synthesized and characterized by melting point determination, solubility, conductivity measurement, elemental analyses, UV-Vis and IR spectroscopy. From the molar conductance value, electronic and infrared spectral data octahedral geometry was proposed for this compound. The iron(II) complex was found to be a more potent antibacterial agent than the ligands and the ciprofloxacin drug. Based on these results, it could be proposed that the novel iron(II) complex can be better

accommodated for pharmaceutical usage as a drug against bacteria infections. From the spectroscopic characterization, it is concluded that ligands act as a neutral bidentate through the HTMA nitrogen atom and carbonyl groups of the succinic acid.

The authors wish to express their sincere thanks to Mr. Bayo Lawal Tudaka, Environmental Consultants Limited, for assisting in the microbial studies. The authors would, also wish to thank Mr. Okoroji of Analtrace Laboratory, for the infrared spectral analysis.

REFERENCES

- Agwara, M. O., Ndifon, P. T., Ndosiri, N. B., Paboudam, A. G., Yufanyi, D. M. & Mohamadou, A. (2010). Synthesis, characterisation and antimicrobial activities of cobalt(II), copper(II) and zinc(II) mixed-ligand complexes containing 1,10-phenanthroline and 2,2'-bipyridine. *Bulletin of the Chemical Society of Ethiopia*, 24(3): 383–389.
- Caires, C.J; Lima, L. S. & Carvalho, C. T. (2010). Thermal behavior of succinic acid, sodium succinate and its compounds with some bivalent transition metal ions in dynamic N_2 and CO_2 atmospheres. *Eclética Quimica*, 35:4-13.
- Caramen M. & Sharaby, M.. (2005). Preparation, characterization and biological activity of Fe(III), Fe(II), Co(II), Ni(II), Cu(II), Zn(II), Cd(II) and $\text{UO}_2(\text{II})$ complexes of new cyclodiphosph(V)azane of

- sulfaguanidine. *Spectrochimica Acta*, 62(3): 326–334.
- Kabbani, A.T., Hammud, H. H. & Ghannoum, A. M. (2007). Preparation and antibacterial activity of copper and cobalt complexes of 4-chloro-3-nitrobenzoate with a nitrogen donor ligand. *Chem Pharm Bull*, 55(3):446-450.
- Kulkarni, A. D., Patil, S. A. and Badami, P. S. (2009). Electrochemical properties of some transition metal complexes: synthesis, characterization and invitro antimicrobial studies of Co(II), Ni(II), Cu(II), Mn(II) and Fe(III) complexes. *International Journal of Electrochemical Science*, 4(5): 717–729.
- Mohammed, S.F; Refat, M.S. & Nashwa, M. (2012). Synthesis of new Series of hexamethylene tetramine complexes with some different metal ions: Spectral, thermal and biological investigations. *Life Science Journal*, 9(2):1243-1253.
- Mohamed, G. G. & Khalil, S. M. (2009). Metal complexes of omeprazole: preparation, spectroscopic and thermal characterization and biological activity. *Journal of Coordination Chemistry*, 62(4): 645–654.
- Nakamoto, K. (2009). *Infrared and Raman Spectra of Inorganic and Coordination Compounds* (6th ed.). John Wiley and Sons: New Jersey, 112-143.
- Patel, R. N., Singh, A., Shukla K. K., Patel, D. K. & Sondhiya, V. P. (2011). Structural, spectroscopic, and biological studies of N, O donor Schiff base copper(II) complexes. *Journal of Coordination Chemistry*, 64(5): 902–919.
- Rai, M., Yadav, A. & Gade, A. (2009). Silver Nanoparticles as a new generation of antimicrobials. *Biotechnology Advances*, 27(1): 76-83.
- Shakru, R., Subhashini, N. J. & Sathish, K. K. (2010). Biological activities of heterocyclic ligands. *Journal of Chemical and Pharmaceutical Research*, 2(1):38-46.
- Zhang, R. Ren, Y. Wang, Q. & Ma, C. (2010). Syntheses and Characterization of 2D and 3D Organotin Polymers with Phenylsuccinic Acid and Trimethyltin Chloride Under Different Conditions. *Journal of Inorganic and Organometallic Polymers and Materials*, 10: 1007-1090.